The Oswestry Disability Index For Low Back Pain

This questionnaire has been designed to give us information as to how your back pain has affected your ability to manage everyday-life activities. Please answer every section, and mark in each section the **one box** that applies to you. Although you may consider that two of the statements in any one section relate to you, please mark the box that **most closely** describes your present-day situation.

| CE. | CTION 4 D | AINI INITENIC | ITV | | | CE/ | TIO. | LC CTAND | INIC | | | | |
|---|--|--|----------------------------------|----------------|-------------|-------|--|--------------------------------|-----------------|----------------|--------------|--------------|--|
| _ | SECTION 1—PAIN INTENSITY My pain is mild to moderate I do not need painkillers | | | | | | SECTION 6—STANDING I can stand as long as I want without extra pain. | | | | | | |
| | My pain is mild to moderate; I do not need painkillers. | | | | | | I can stand as long as I want without extra pain. | | | | | | |
| | - | The pain is bad, but I manage without taking painkillers. Painkillers give complete relief from pain. | | | | | I can stand as long as I want, but it gives me extra pain. Pain prevents me from standing for more than 1 hour. | | | | | | |
| | | | | - | | | | i prevents me i prevents me | | | | • | |
| | | _ | ite relief from | | | | | | | | | | |
| | | - | le relief from ect on the pai | - | | | | prevents me | | | an 10 minute | 5. | |
| Ш | Pallikillers | nave no ene | ct on the pai | 11. | | | Pall | prevents me | e iroiii staiiu | ilig at all. | | | |
| SEC | SECTION 2—PERSONAL CARE | | | | | | | SECTION 7—SLEEPING | | | | | |
| | I can look a | after myself | normally with | nout causing | extra pain. | | Pair | does not pr | event me fro | m sleeping v | well. | | |
| | I can look a | after myself | normally, but | t it causes ex | tra pain. | | I sle | ep well, but | only when ta | king medica | tion. | | |
| | It is painfu | l to look afte | r myself, and | l I am slow ar | nd careful. | | Eve | n when I take | e medication, | , I sleep less | than 6 hours | i . | |
| | I need som | need some help but manage most of my personal care. | | | | | Even when I take medication, I sleep less than 4 hours. | | | | | | |
| | | I need help every day in most aspects of self-care. | | | | | | n when I take | | | than 2 hours | . | |
| | I do not ge | t dressed. I v | wash with dif | ficulty and st | ay in bed. | | Pair | prevents me | e from sleepi | ng at all. | | | |
| SEC | CTION 3—LI | FTING | | | | SEC | SECTION 8—SOCIAL LIFE | | | | | | |
| | I can lift heavy weights without causing extra pain. | | | | | | ☐ My social life is normal and causes me no extra pain. | | | | | | |
| | | I can lift heavy weights, but it gives me extra pain. | | | | | My social life is normal, but increases the degree of pain. | | | | | | |
| | Pain prevents me from lifting heavy weights off the floor, | | | | | | Pain affects my social life by limiting only my more energetic | | | | | | |
| | but I can manage if items are conveniently positioned, i.e. | | | | | | interests, such as dancing, sports, etc. | | | | | | |
| | on a table. | | | | | | Pain affects my social life, and I do not go out as often. | | | | | | |
| | Pain preve | Pain prevents me from lifting heavy weights, but I can | | | | | Pain has restricted my social life to my home. | | | | | | |
| | manage light weights if they are conveniently positioned. | | | | | | I have no social life because of pain. | | | | | | |
| | I can lift only very light weights. | | | | | | | | | • | | | |
| | I cannot lift or carry anything at all. | | | | | | | SECTION 9—TRAVELING | | | | | |
| | | | | | | | ☐ I can travel anywhere without extra pain.☐ I can travel anywhere, but it gives me extra pain. | | | | | | |
| SECTION 4—WALKING | | | | | | | I can travel anywhere, but it gives me extra pain. | | | | | | |
| | I can walk as far as I wish. | | | | | | Pain is bad, but I manage journeys over 2 hours. | | | | | | |
| | Pain prevents me from walking more than 1 mile. | | | | | | Pain restricts me to journeys of less than 1 hour. | | | | | | |
| | Pain prevents me from walking more than 1/2 mile. | | | | | | Pain restricts me to necessary journeys under 1/2 hour. | | | | | | |
| | Pain prevents me from walking more than 1/4 mile. I can walk only if I use a cane or crutches. | | | | | | ☐ Pain prevents traveling except to the doctor/hospital. | | | | | | |
| | | - | | | | SEC | TION | I 10—CHAN | GING DEGRI | FF OF PAIN | | | |
| ☐ I am in bed or in a chair for most of every day. | | | | | | | SECTION 10—CHANGING DEGREE OF PAIN My pain is rapidly getting better. | | | | | | |
| SECTION 5—SITTING | | | | | | _ | My pain braphary getting better.My pain fluctuates but overall is definitely getting better. | | | | | | |
| | | | | | | | My pain seems to be getting better but improvement is slow at | | | | | | |
| | | I can sit in my favorite chair only, but for as long as I like. | | | | | | present. | | | | | |
| | Pain prevents me from sitting for more than 1 hour. | | | | | | ☐ My pain is neither getting better nor worse. | | | | | | |
| | Pain prevents me from sitting for more than 1/2 hour. | | | | | | _ | | | | | | |
| | Pain prevents me from sitting for more than 10 minutes. | | | | | | | | | | | | |
| □ Pain prevents me from sitting at all. | | | | | | | | | | | | | |
| г. | ain Cararit | Scalor | | | | | | | | | | | |
| Pain Severity Scale: Rate your usual level of pain today by checking one box on the following scale. | | | | | | | | | | | | | |
| | ate you | ı usuai il | -vei oi þ | iii touay | , by check | viiig | UITE | DOX OII | LITE TOTIO | Talle Sca | | | |
| | 0 | 1 | 2 | 3 | 4 | 5 | | 6 | 7 | 8 | 9 | 10 | |
| L | No Pain | | | <u> </u> | | | | · · · | <u>'</u> | | 1 - | Excruciating | |
| | • | | | | | | | | | | | Pain | |

DATE:

SCORE:

Office use only

PATIENT NAME:_